

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/EP05/002563

International filing date: 10 February 2005 (10.02.2005)

Document type: Certified copy of priority document

Document details: Country/Office: US  
Number: 60/637,120  
Filing date: 17 December 2004 (17.12.2004)

Date of receipt at the International Bureau: 31 March 2005 (31.03.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

1:0 FEB 2005

PA 1273841



# THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office


January 19, 2005

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/637,120

FILING DATE: December 17, 2004

By Authority of the  
COMMISSIONER OF PATENTS AND TRADEMARKS



T. LAWRENCE  
Certifying Officer

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

**PROVISIONAL APPLICATION FOR PATENT COVER SHEET**

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No. ED39888246US**INVENTOR(S)**

Given Name (first and middle (if any))	Family Name or Surname	Residence (City and either State or Foreign Country)
Jean-Marie	Gouot	69450 St. Cyr Au Mont D'Or, France
Marie-Claire	Grosjean-Cournoyer	69250 Curis Au Mont D'Or, France

Additional inventors are being named on the \_\_\_\_\_ separately numbered sheets attached hereto

**TITLE OF THE INVENTION (500 characters max):**

FUNGICIDAL COMPOSITION COMPRISING A PYRIDYLETHYLBENZAMIDE DERIVATIVE AND COMPOUND CAPABLE OF INHIBITING THE TRANSPORT OF ELECTRONS OF THE RESPIRATORY CHAIN IN PHYTOPATHOGENIC FUNGAL ORGANISMS

Direct all correspondence to:

**CORRESPONDENCE ADDRESS**☒ The address corresponding to Customer Number:

34469

OR

☐ Firm or  
Individual Name

Address

City

State

Zip

Country

Telephone

Fax

**ENCLOSED APPLICATION PARTS (check all that apply)**☐ Application Data Sheet. See 37 CFR 1.76☐ CD(s), Number of CDs \_\_\_\_\_☒ Specification Number of Pages \_\_\_\_\_☐ Other (specify) \_\_\_\_\_☐ Drawing(s) Number of Sheets \_\_\_\_\_**Application Size Fee:** If the specification and drawings exceed 100 sheets of paper, the application size fee due is \$250 (\$125 for small entity) for each additional 50 sheets or fraction thereof. See 35 U.S.C. 41(a)(1)(G) and 37 CFR 1.16(s).**METHOD OF PAYMENT OF FILING FEES AND APPLICATION SIZE FEE FOR THIS PROVISIONAL APPLICATION FOR PATENT**☐ Applicant claims small entity status. See 37 CFR 1.27.

TOTAL FEE AMOUNT (\$)

☐ A check or money order is enclosed to cover the filing fee and application size fee (if applicable).

200.00

☐ Payment by credit card. Form PTO-2038 is attached☒ The Director is hereby authorized to charge the filing fee and application size fee (if applicable) or credit any overpayment to DepositAccount Number: 502510

A duplicative copy of this form is enclosed for fee processing.

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☒ No.☐ Yes, the name of the U.S. Government agency and the Government contract number are: \_\_\_\_\_

SIGNATURE

Richard E.L. HendersonDate 17 December 2004TYPED or PRINTED NAME Richard E.L. HendersonREGISTRATION NO. 31,619

(if appropriate)

TELEPHONE (412) 777-3916Docket Number: CS8430/BCS044015**USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT**

This collection of information is required by 37 CFR 1.51. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.11 and 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

Effective on 12/08/2004.

Fees pursuant to the Consolidated Appropriations Act, 2005 (H.R. 4818).

**FEE TRANSMITTAL**  
**For FY 2005**☐ Applicant claims small entity status. See 37 CFR 1.27TOTAL AMOUNT OF PAYMENT (\$)  
200.00**Complete if Known**

Application Number	To be Assigned
Filing Date	Herewith
First Named Inventor	Jean Marie Gouot et al
Examiner Name	
Art Unit	
Attorney Docket No.	PO8430/BCS044015

**METHOD OF PAYMENT (check all that apply)**
☐ Check ☐ Credit Card ☐ Money Order ☐ None ☐ Other (please identify): \_\_\_\_\_

☒ Deposit Account Deposit Account Number: 502510 Deposit Account Name: Bayer CropScience LP

For the above-identified deposit account, the Director is hereby authorized to: (check all that apply)

☒ Charge fee(s) indicated below ☐ Charge fee(s) indicated below, except for the filing fee

☒ Charge any additional fee(s) or underpayments of fee(s) under 37 CFR 1.16 and 1.17 ☒ Credit any overpayments

WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.

**FEE CALCULATION****1. BASIC FILING, SEARCH, AND EXAMINATION FEES**

Application Type	FILING FEES		SEARCH FEES		EXAMINATION FEES		Fees Paid (\$)
	Fee (\$)	Small Entity Fee (\$)	Fee (\$)	Small Entity Fee (\$)	Fee (\$)	Small Entity Fee (\$)	
Utility	300	150	500	250	200	100	
Design	200	100	100	50	130	65	
Plant	200	100	300	150	160	80	
Reissue	300	150	500	250	600	300	
Provisional	200	100	0	0	0	0	200.00

**2. EXCESS CLAIM FEES****Fee Description**

	Fee (\$)	Small Entity Fee (\$)
Each claim over 20 or, for Reissues, each claim over 20 and more than in the original patent	50	25
Each independent claim over 3 or, for Reissues, each independent claim more than in the original patent	200	100
Multiple dependent claims	360	180

<b>Total Claims</b>	<b>Extra Claims</b>	<b>Fee (\$)</b>	<b>Fee Paid (\$)</b>	<b>Multiple Dependent Claims</b>	<b>Fee (\$)</b>	<b>Fee Paid (\$)</b>
- 20 or HP = _____ x _____ = _____						
HP = highest number of total claims paid for, if greater than 20						
<b>Indep. Claims</b>	<b>Extra Claims</b>	<b>Fee (\$)</b>	<b>Fee Paid (\$)</b>			
- 3 or HP = _____ x _____ = _____						
HP = highest number of independent claims paid for, if greater than 3						

**3. APPLICATION SIZE FEE**

If the specification and drawings exceed 100 sheets of paper, the application size fee due is \$250 (\$125 for small entity) for each additional 50 sheets or fraction thereof. See 35 U.S.C. 41(a)(1)(G) and 37 CFR 1.16(s).

<b>Total Sheets</b>	<b>Extra Sheets</b>	<b>Number of each additional 50 or fraction thereof</b>	<b>Fee (\$)</b>	<b>Fee Paid (\$)</b>
23	-77	/ 50 = _____ (round up to a whole number) x	250.00	= 0.00

**4. OTHER FEE(S)**

Non-English Specification, \$130 fee (no small entity discount)

Other: \_\_\_\_\_

**SUBMITTED BY**

Signature	<i>Richard E.L. Henderson</i>	Registration No.	31619	Telephone	412-777-3916
Name (Print/Type)	Richard E.L. Henderson	(Attorney/Agent)		Date	December 17, 2004

This collection of information is required by 37 CFR 1.136. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 30 minutes to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

**Fungicidal composition comprising a pyridylethylbenzamide derivative and a compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms**


5 The present invention relates to novel fungicide compositions comprising a pyridylethylbenzamide derivative and a compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms. The present invention also relates to a method of combating or controlling phytopathogenic fungi by applying at a locus infested or liable to be infested such a composition.

International patent application WO 01/11965 generically discloses numerous pyridylethylbenzamide derivatives. The possibility of combining one or more of these numerous pyridylethylbenzamide derivatives with known fungicidal products to develop a fungicidal activity is disclosed in general terms, without any specific example or biological data.

It is always of high-interest in agriculture to use novel pesticidal mixtures showing a synergistic effect in order notably to avoid or to control the development of resistant strains to the active ingredients or to the mixtures of known active ingredients used by the farmer while minimising the doses of chemical products spread in the environment and reducing the cost of the treatment.

We have now found some novel fungicidal compositions which possess the above mentioned characteristics.

Accordingly, the present invention relates to a composition comprising :  
a) a pyridylethylbenzamide derivative of general formula (I)



in which :

- p is an integer equal to 1, 2, 3 or 4;
- q is an integer equal to 1, 2, 3, 4 or 5;
- each substituent X is chosen, independently of the others, as being halogen, alkyl or haloalkyl;

- each substituent Y is chosen, independently of the others, as being halogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxy, amino, phenoxy, alkylthio, dialkylamino, acyl, cyano, ester, hydroxy, aminoalkyl, benzyl, haloalkoxy, halosulphonyl, halothioalkyl, alkoxyalkenyl, alkylsulphonamide, nitro, alkylsulphonyl, phenylsulphonyl or benzylsulphonyl;

as to the N-oxides of the compounds thereof;  
and

b) a compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms;

in a (a) / (b) weight ratio of from 0.01 to 20.

In the context of the present invention, a compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms is chosen from the group comprising a compound capable of inhibiting reduced nicotinamide-adenine dinucleotide (NADH) deshydrogenase in phytopathogenic fungal organisms, a compound capable of inhibiting succinate dehydrogenase in phytopathogenic fungal organisms and a compound capable of inhibiting mitochondrial ubiquinol:ferricytochrome-c oxidoreductase in phytopathogenic fungal organisms.

A compound capable of inhibiting NADH deshydrogenase in phytopathogenic fungal organisms is also known as complex I inhibitor.

A compound capable of succinate dehydrogenase in phytopathogenic fungal organisms is also known as complex II inhibitor.

A compound capable of inhibiting mitochondrial ubiquinol:ferricytochrome-c oxidoreductase in phytopathogenic fungal organisms is also known as complex III inhibitor.

In the context of the present invention :

- halogen means chlorine, bromine, iodine or fluorine;

- each of the alkyl or acyl radicals present in the molecule contains from 1 to 10 carbon atoms, preferably from 1 to 7 carbon atoms, more preferably from 1 to 5 carbon atoms, and may be linear or branched;

- each of the alkenyl or alkynyl radicals present in the molecule contains from 2 to 10 carbon atoms, preferably from 2 to 7 carbon atoms, more preferably from 2 to 5

carbon atoms, and may be linear or branched.

The composition according to the present invention provides a synergistic effect. This synergistic effect allows a reduction of the chemical substances spread into the environment and a reduction of the cost of the fungal treatment.

In the context of the present invention, the term "synergistic effect" is defined by Colby according to the article entitled "Calculation of the synergistic and antagonistic responses of herbicide combinations" Weeds, (1967), 15, pages 20-22.

The latter article mentions the formula:

$$E = x + y - \frac{x * y}{100}$$

in which E represents the expected percentage of inhibition of the disease for the combination of the two fungicides at defined doses (for example equal to x and y respectively), x is the percentage of inhibition observed for the disease by the compound (I) at a defined dose (equal to x), y is the percentage of inhibition observed for the disease by the compound (II) at a defined dose (equal to y). When the percentage of inhibition observed for the combination is greater than E, there is a synergistic effect.

The composition according to the present invention comprises a pyridylethylbenzamide derivative of general formula (I). Preferably, the present invention relates to a composition comprising a pyridylethylbenzamide derivative of general formula (I) in which the different characteristics may be chosen alone or in combination as being :

- as regards p, p is 2;
- as regards q, q is 1 or 2. More preferably, q is 2;
- as regards X, X is chosen, independently of the others, as being halogen or haloalkyl. More preferably, X is chosen, independently of the others, as being a chlorine atom or a trifluoromethyl group;
- as regards Y, Y is chosen, independently of the others, as being halogen or haloalkyl. More preferably, Y is chosen, independently of the others, as being a chlorine atom or a trifluoromethyl group;

More preferably, the pyridylethylbenzamide derivative of general formula (I) present in the composition of the present invention is :

- N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (compound 1);
- N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-iodobenzamide (compound 2); or

- N-{2-[3,5-dichloro-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (compound 3).

Even more preferably, the pyridylethylbenzamide derivative of general formula (I) present in the composition of the present invention is N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (compound 1).

5

The composition according to the present invention comprises a compound capable of inhibiting the transport of electrons of the respiratory chain.

Preferably, the present invention also relates to a composition comprising a complex I inhibitor. More preferably, the present invention relates to a composition comprising a complex I inhibitor which is diflumeritin.

Preferably, the present invention also relates to a composition comprising a complex II inhibitor. More preferably, the present invention relates to a composition comprising a complex II inhibitor selected from the carboxamide derivatives which may for example be *N*-[2-(1,3-dimethyl-butyl)-phenyl]-5-fluoro-1,3-dimethyl-1*H*-pyrazole-4-carboxamide, *N*-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoromethyl)-1-methyl-1*H*-pyrazole-4-carboxamide, *N*-[2-(1,3-dimethylbutyl)-thiophen-3-yl] 1-methyl-3-(trifluoromethyl)-1*H*-pyrazole-4-carboxamide, benodanil, carboxin, fenfuram, flutolanil, furametpyr, mepronil, boscalid, oxycarboxin, thifluzamide. *N*-[2-(1,3-dimethyl-butyl)-phenyl]-5-fluoro-1,3-dimethyl-1*H*-pyrazole-4-carboxamide, *N*-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoro-methyl)-1-methyl-1*H*-pyrazole-4-carboxamide, boscalid and oxycarboxin are preferred.

Preferably, the present invention also relates to a composition comprising a complex III inhibitor. More preferably, the present invention relates to a composition comprising a complex III inhibitor selected from strobilurin derivatives, cyazofamid, fenamidone or famoxadone. Fenamidone is preferred.

Strobilurin derivatives are also preferred. According to the present invention, strobilurin derivatives may for example be azoxystrobin, dimoxystrobin, fluoxastrobin, kresoxim-methyl, metominostrobin, trifloxystrobin, pyraclostrobin, picoxystrobin or 2-{2-[6-(3-chloro-2-methylphenoxy)-5-fluoro-pyrimidin-4-yloxy]-phenyl}2-methoxyimino-*N*-methylacetamide. 2-{2-[6-(3-chloro-2-methylphenoxy)-5-fluoro-pyrimidin-4-yloxy]-phenyl}2-methoxyimino-*N*-methylacetamide, azoxystrobin, trifloxystrobin and fluoxastrobin are preferred.



The composition according to the present invention comprises (a) at least a pyridylethylbenzamide derivative of general formula (I) and (b) a compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms in an (a) / (b) weight ratio of from 0.01 to 20; preferably of from 0.05 to 10; even more preferably, of from 0.1 to 5.

The composition of the present invention may further comprise at least one other fungicide active ingredient (c).

The fungicidal active ingredient (c) may be selected from azaconazole, azoxystrobin, (Z)-N-[ $\alpha$ -(cyclopropylmethoxyimino)-2,3-difluoro-6-(trifluoromethyl)benzyl]-2-phenylacetamide, benalaxyl, benomyl, benthiavalicarb, biphenyl, bitertanol, blasticidin-S, boscalid, borax, bromuconazole, bupirimate, sec-butylamine, calcium polysulfide, captafol, captan, carbendazim, carboxin, carpropamid, chinomethionat, chlorothalonil, chlozolate, copper hydroxide, copper octanoate, copper oxychloride, copper sulfate, cuprous oxide, cymoxanil, cyproconazole, cyprodinil, dazomet, debacarb, dichlofluanid, dichlorophen, diclocymet, diclomezine, dicloran, diethofencarb, difenoconazole, difenzoquat metilsulfate, difenzoquat, diflumetorim, dimethirimol, dimethomorph, diniconazole, dinobuton, dinocap, diphenylamine, dithianon, dodemorph, dodemorph acetate, dodine, edifenphos, epoxiconazole, ethaboxam, ethirimol, ethoxyquin, etridiazole, famoxadone, fenamidone, fenarimol, fenbuconazole, fenfuram, fenhexamid, fempiclonil, fenoxanil, fenpropidin, fenpropimorph, fentin, fentin hydroxide, fentin acetate, ferbam, ferimzone, fluazinam, fludioxonil, fluoroimide, fluoxastrobin, fluquinconazole, flusilazole, flusulfamide, flutolanil, flutriafol, folpet, formaldehyde, fosetyl, fosetyl-aluminium, fuberidazole, furalaxyl, furametpyr, guazatine, guazatine acetates, hexachlorobenzene, hexaconazole, 8-hydroxyquinoline sulfate, potassium hydroxyquinoline sulfate, hymexazol, cyazofamid, imazalil sulfate, imazalil, imibenconazole, iminocadine, iminocadine triacetate, ipconazole, iprobenfos, iprodione, iprovalicarb, isoprothiolane, kasugamycin, kasugamycin hydrochloride hydrate, kresoxim-methyl, mancozeb, maneb, mepanipyrim, mepronil, mercuric chloride, mercuric oxide, mercurous chloride, metalaxyl, metalaxyl-M, metam-sodium, metam, metconazole, methasulfocarb, methyl isothiocyanate, metiram, metominostrobin, mildiomyacin, myclobutanil, nabam, nickel bis(dimethyldithiocarbamate), nitrothal-isopropyl, nuarimol, octhilinone, ofurace, oleic acid, oxadixyl, oxine-copper, oxpoconazole fumarate, oxycarboxin, pefurazate, penconazole, pencycuron, pentachlorophenol, sodium

pentachlorophenoxide, pentachlorophenyl laurate, phenylmercury acetate, sodium 2-phenylphenoxide, 2-phenylphenol, phthalide, picoxystrobin, piperalin, polyoxins, polyoxin B, polyoxin, polyoxorim, probenazole, prochloraz, procymidone, propamocarb hydrochloride, propamocarb, propiconazole, propineb, 5 prothioconazole, pyrazophos, pyributicarb, pyrifenox, pyrimethanil, pyroquilon, quinoxifen, quintozone, silthiofam, spiroxamine, sulfur, tar oils, tebuconazole, tecnazene, tetraconazole, thiabendazole, thifluzamide, thiophanate-methyl, thiram, tolclofos-methyl, tolylfluanid, triadimefon, triadimenol, triazoxide, tricyclazole, tridemorph, trifloxystrobin, triflumizole, triforine, triticonazole, validamycin, 10 vinclozolin, zineb, ziram, zoxamide, phosphorous acid, pyraclostrobin and simeconazole.

Preferably, fungicidal active ingredient (c) is selected from captane, folpet, dodine, propineb, mancozeb, thiram, tolylfluanid, iminoctadine, dithianon, copper hydroxide, copper octanoate, copper oxychloride, copper sulfate, fosetyl-Al, 15 phosphorous acid, cymoxanil, iprovalicarb, bentiavalicarb, chlorotalonil, propamocarb, tebuconazole and spiroxamine.

Where the third active ingredient (c) as defined above is present in the composition, this compound may be present in an amount of (a) : (b) : (c) weight 20 ratio of from 1 : 0.01 : 0.01 to 1 : 20 : 20; the ratios of compound (a) and compound (c) varying independently from each other. Preferably, the (a) : (b) : (c) weight ratio may be of from 1 : 0.05 : 0.05 to 1 : 10 : 10.

Following compositions may be cited to illustrate in a non-limited manner the 25 present invention : compound 1 with diflometorin, compound 1 with *N*-[2-(1,3-dimethyl-butyl)-phenyl]-5-fluoro-1,3-dimethyl-1*H*-pyrazole-4-carboxamide, compound 1 with *N*-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoro-methyl)-1-methyl-1*H*-pyrazole-4-carboxamide, compound 1 with *N*-[2-(1,3-dimethylbutyl)-thiophen-3-yl] 1-methyl-3-(trifluoromethyl)-1*H*-pyrazole-4-carboxamide, compound 30 1 with benodanil, compound 1 with carboxin, compound 1 with fenfuram, compound 1 with flutolanil, compound 1 with furametpyr, compound 1 with mepronil, compound 1 with boscalid, compound 1 with oxycarboxin, compound 1 with thifluzamide, compound 1 with cyazofamid, compound 1 with fenamidone, compound 1 with famoxadone, azoxystrobin, compound 1 with dimoxystrobin, 35 compound 1 with fluoxastrobin, compound 1 with kresoxim-methyl, compound 1 with metominostrobin, compound 1 with trifloxystrobin, compound 1 with

pyraclostrobin, compound 1 with picoxystrobin, compound 1 with 2-{2-[6-(3-chloro-2-methylphenoxy)-5-fluoro-pyrimidin-4-yloxy]-phenyl}2-methoxyimino-N-methylacetamide, compound 2 with diflumetorin, compound 2 with *N*-[2-(1,3-dimethyl-butyl)-phenyl]-5-fluoro-1,3-dimethyl-1*H*-pyrazole-4-carboxamide,

5 compound 2 with *N*-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoro-methyl)-1-methyl-1*H*-pyrazole-4-carboxamide, compound 2 with *N*-[2-(1,3-dimethylbutyl)-thiophen-3-yl] 1-methyl-3-(trifluoromethyl)-1*H*-pyrazole-4-carboxamide, compound 2 with benodanil, compound 2 with carboxin, compound 2 with fenfuram, compound 2 with flutolanil, compound 2 with furametpyr, compound 2 with mepronil,

10 compound 2 with boscalid, compound 2 with oxycarboxin, compound 2 with thifluzamide, compound 2 with cyazofamid, compound 2 with fenamidone, compound 2 with famoxadone. azoxystrobin, compound 2 with dimoxystrobin, compound 2 with fluoxastrobin, compound 2 with kresoxim-methyl, compound 2 with metominostrobin, compound 2 with trifloxystrobin, compound 2 with

15 pyraclostrobin, compound 2 with picoxystrobin, compound 2 with 2-{2-[6-(3-chloro-2-methylphenoxy)-5-fluoro-pyrimidin-4-yloxy]-phenyl}2-methoxyimino-N-methylacetamide, compound 3 with diflumetorin, compound 3 with *N*-[2-(1,3-dimethyl-butyl)-phenyl]-5-fluoro-1,3-dimethyl-1*H*-pyrazole-4-carboxamide,

20 compound 3 with *N*-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoro-methyl)-1-methyl-1*H*-pyrazole-4-carboxamide, compound 3 with *N*-[2-(1,3-dimethylbutyl)-thiophen-3-yl] 1-methyl-3-(trifluoromethyl)-1*H*-pyrazole-4-carboxamide, compound 3 with benodanil, compound 3 with carboxin, compound 3 with fenfuram, compound 3 with flutolanil, compound 3 with furametpyr, compound 3 with mepronil,

25 compound 3 with boscalid, compound 3 with oxycarboxin, compound 3 with thifluzamide, compound 3 with cyazofamid, compound 3 with fenamidone, compound 3 with famoxadone. azoxystrobin, compound 3 with dimoxystrobin, compound 3 with fluoxastrobin, compound 3 with kresoxim-methyl, compound 3 with metominostrobin, compound 3 with trifloxystrobin, compound 3 with

30 pyraclostrobin, compound 3 with picoxystrobin, compound 3 with 2-{2-[6-(3-chloro-2-methylphenoxy)-5-fluoro-pyrimidin-4-yloxy]-phenyl}2-methoxyimino-N-methylacetamide.

The composition according to the present invention may further comprise an other additional component such as an agriculturally acceptable support, carrier or

35 filler.

In the present specification, the term "support" denotes a natural or synthetic, organic or inorganic material with which the active material is combined to make it easier to apply, notably to the parts of the plant. This support is thus generally inert and should be agriculturally acceptable. The support may be a solid or a liquid.

5 Examples of suitable supports include clays, natural or synthetic silicates, silica, resins, waxes, solid fertilisers, water, alcohols, in particular butanol, organic solvents, mineral and plant oils and derivatives thereof. Mixtures of such supports may also be used.

The composition may also comprise other additional components. In particular, the composition may further comprise a surfactant. The surfactant can be an emulsifier, a dispersing agent or a wetting agent of ionic or non-ionic type or a mixture of such surfactants. Mention may be made, for example, of polyacrylic acid salts, lignosulphonic acid salts, phenolsulphonic or naphthalenesulphonic acid salts, polycondensates of ethylene oxide with fatty alcohols or with fatty acids or with fatty amines, substituted phenols (in particular alkylphenols or arylphenols), salts of sulphosuccinic acid esters, taurine derivatives (in particular alkyl taurates), phosphoric esters of polyoxyethylated alcohols or phenols, fatty acid esters of polyols, and derivatives of the above compounds containing sulphate, sulphonate and phosphate functions. The presence of at least one surfactant is generally essential

15 when the active material and/or the inert support are water-insoluble and when the vector agent for the application is water. Preferably, surfactant content may be comprised between 5% and 40% by weight of the composition.

Additional components may also be included, e.g. protective colloids, adhesives, thickeners, thixotropic agents, penetration agents, stabilisers, sequestering agents. More generally, the active materials can be combined with any solid or liquid additive, which complies with the usual formulation techniques.

25

In general, the composition according to the invention may contain from 0.05 to 99% (by weight) of active material, preferably 10 to 70% by weight.

Compositions according to the present invention can be used in various forms such as aerosol dispenser, capsule suspension, cold fogging concentrate, dustable powder, emulsifiable concentrate, emulsion oil in water, emulsion water in oil, encapsulated granule, fine granule, flowable concentrate for seed treatment, gas (under pressure), gas generating product, granule, hot fogging concentrate, macrogranule, microgranule, oil dispersible powder, oil miscible flowable concentrate, oil miscible liquid, paste, plant rodlet, powder for dry seed treatment,

30 seed coated with a pesticide, soluble concentrate, soluble powder, solution for seed

treatment, suspension concentrate (flowable concentrate), ultra low volume (ulv) liquid, ultra low volume (ulv) suspension, water dispersible granules or tablets, water dispersible powder for slurry treatment, water soluble granules or tablets, water soluble powder for seed treatment and wettable powder.

5        These compositions include not only compositions which are ready to be applied to the plant or seed to be treated by means of a suitable device, such as a spraying or dusting device, but also concentrated commercial compositions which must be diluted before they are applied to the crop.

10        The fungicidal compositions of the present invention can be used to curatively or preventively control phytopathogenic fungi of crops. Thus, according to a further aspect of the present invention, there is provided a method for preventively or curatively controlling phytopathogenic fungi of crops characterised in that a fungicidal composition as hereinbefore defined is applied to the seed, the plant and/or to the fruit  
15 of the plant or to the soil in which the plant is growing or in which it is desired to grow.

The composition as used against phytopathogenic fungi of crops comprises an effective and non-phytotoxic amount of an active material of general formula (I).

20        The expression "effective and non-phytotoxic amount" means an amount of composition according to the invention which is sufficient to control or destroy the fungi present or liable to appear on the crops, and which does not entail any appreciable symptom of phytotoxicity for the said crops. Such an amount can vary within a wide range depending on the fungus to be combated or controlled, the type of crop, the climatic conditions and the compounds included in the fungicidal composition according to the invention.

25        This amount can be determined by systematic field trials, which are within the capabilities of a person skilled in the art.

30        The method of treatment according to the present invention is useful to treat propagation material such as tubers or rhizomes, but also seeds, seedlings or seedlings pricking out and plants or plants pricking out. This method of treatment can also be useful to treat roots. The method of treatment according to the present invention can also be useful to treat the overground parts of the plant such as trunks, stems or stalks, leaves, flowers and fruits of the concerned plant.

35        Among the plants that can be protected by the method according to the invention, mention may be made of cotton; flax; vine; fruit crops such as *Rosaceae* sp. (for instance pip fruits such as apples and pears, but also stone fruits such as apricots, almonds and peaches), *Ribesioideae* sp., *Juglandaceae* sp., *Betulaceae* sp.,

*Anacardiaceae sp.*, *Fagaceae sp.*, *Moraceae sp.*, *Oleaceae sp.*, *Actinidaceae sp.*, *Lauraceae sp.*, *Musaceae sp.* (for instance banana trees and plantains), *Rubiaceae sp.*, *Theaceae sp.*, *Sterculiaceae sp.*, *Rutaceae sp.* (for instance lemons, oranges and grapefruits); leguminous crops such as *Solanaceae sp.* (for instance tomatoes),  
 5 *Liliaceae sp.*, *Asteraceae sp.* (for instance lettuces), *Umbelliferae sp.*, *Cruciferae sp.*, *Chenopodiaceae sp.*, *Cucurbitaceae sp.*, *Papilionaceae sp.* (for instance peas), *Rosaceae sp.* (for instance strawberries); big crops such as *Graminae sp.* (for instance maize, cereals such as wheat, rice, barley and triticales), *Asteraceae sp.* (for instance sunflower), *Cruciferae sp.* (for instance colza), *Papilionaceae sp.* (for  
 10 instance soja), *Solanaceae sp.* (for instance potatoes), *Chenopodiaceae sp.* (for instance beetroots); horticultural and forest crops; as well as genetically modified homologues of these crops.

Among the plants and the possible diseases of these plants protected by the method according to the present invention, mention may be made of :

- 15 - wheat, as regards controlling the following seed diseases: fusaria (*Microdochium nivale* and *Fusarium roseum*), stinking smut (*Tilletia caries*, *Tilletia controversa* or *Tilletia indica*), septoria disease (*Septoria nodorum*) and loose smut;
- wheat, as regards controlling the following diseases of the aerial parts of the plant: cereal eyespot (*Tapesia yellundae*, *Tapesia acuiformis*), take-all  
 20 (*Gaeumannomyces graminis*), foot blight (*F. culmorum*, *F. graminearum*), black speck (*Rhizoctonia cerealis*), powdery mildew (*Erysiphe graminis forma specie tritici*), rusts (*Puccinia striiformis* and *Puccinia recondita*) and septoria diseases (*Septoria tritici* and *Septoria nodorum*);
- wheat and barley, as regards controlling bacterial and viral diseases, for  
 25 example barley yellow mosaic;
- barley, as regards controlling the following seed diseases: net blotch (*Pyrenophora graminea*, *Pyrenophora teres* and *Cochliobolus sativus*), loose smut (*Ustilago nuda*) and fusaria (*Microdochium nivale* and *Fusarium roseum*);
- barley, as regards controlling the following diseases of the aerial parts of the  
 30 plant: cereal eyespot (*Tapesia yellundae*), net blotch (*Pyrenophora teres* and *Cochliobolus sativus*), powdery mildew (*Erysiphe graminis forma specie hordei*), dwarf leaf rust (*Puccinia hordei*) and leaf blotch (*Rhynchosporium secalis*);
- potato, as regards controlling tuber diseases (in particular *Helminthosporium solani*, *Phoma tuberosa*, *Rhizoctonia solani*, *Fusarium solani*), mildew (*Phytophthora*  
 35 *infestans*) and certain viruses (virus Y);

- potato, as regards controlling the following foliage diseases: early blight (*Alternaria solani*), mildew (*Phytophthora infestans*);

- cotton, as regards controlling the following diseases of young plants grown from seeds: damping-off and collar rot (*Rhizoctonia solani*, *Fusarium oxysporum*) and  
5 black root rot (*Thielaviopsis basicola*);

- protein yielding crops, for example peas, as regards controlling the following seed diseases: anthracnose (*Ascochyta pisi*, *Mycosphaerella pinodes*), fusaria (*Fusarium oxysporum*), grey mould (*Botrytis cinerea*) and mildew (*Peronospora pisi*);

10 - oil-bearing crops, for example rape, as regards controlling the following seed diseases: *Phoma lingam*, *Alternaria brassicae* and *Sclerotinia sclerotiorum*;

- corn, as regards controlling seed diseases: (*Rhizopus* sp., *Penicillium* sp., *Trichoderma* sp., *Aspergillus* sp., and *Gibberella fujikuroi*);

- flax, as regards controlling the seed disease: *Alternaria linicola*;

15 - forest trees, as regards controlling damping-off (*Fusarium oxysporum*, *Rhizoctonia solani*);

- rice, as regards controlling the following diseases of the aerial parts: blast disease (*Magnaporthe grisea*), bordered sheath spot (*Rhizoctonia solani*);

- leguminous crops, as regards controlling the following diseases of seeds or  
20 of young plants grown from seeds: damping-off and collar rot (*Fusarium oxysporum*, *Fusarium roseum*, *Rhizoctonia solani*, *Pythium* sp.);

- leguminous crops, as regards controlling the following diseases of the aerial parts: grey mould (*Botrytis* sp.), powdery mildews (in particular *Erysiphe cichoracearum*, *Sphaerotheca fuliginea* and *Leveillula taurica*), fusaria (*Fusarium oxysporum*, *Fusarium roseum*), leaf spot (*Cladosporium* sp.), alternaria leaf spot  
25 (*Alternaria* sp.), anthracnose (*Colletotrichum* sp.), septoria leaf spot (*Septoria* sp.), black speck (*Rhizoctonia solani*), mildews (for example *Bremia lactucae*, *Peronospora* sp., *Pseudoperonospora* sp., *Phytophthora* sp.);

- fruit trees, as regards diseases of the aerial parts: monilia disease (*Monilia fructigenae*, *M. laxa*), scab (*Venturia inaequalis*), powdery mildew (*Podosphaera leucotricha*);  
30

- vine, as regards diseases of the foliage: in particular grey mould (*Botrytis cinerea*), powdery mildew (*Uncinula necator*), black rot (*Guignardia biwelli*) and mildew (*Plasmopara viticola*);

- beetroot, as regards the following diseases of the aerial parts: cercospora blight (*Cercospora beticola*), powdery mildew (*Erysiphe beticola*), leaf spot (*Ramularia beticola*).

5           The fungicidal composition according to the present invention may also be used against fungal diseases liable to grow on or inside timber. The term "timber" means all types of species of wood, and all types of working of this wood intended for construction, for example solid wood, high-density wood, laminated wood, and plywood. The method for treating timber according to the invention mainly consists  
10 in contacting one or more compounds of the present invention, or a composition according to the invention; this includes for example direct application, spraying, dipping, injection or any other suitable means.

          The fungicidal composition according to the present invention may also be  
15 used in the treatment of genetically modified organisms with the compounds according to the invention or the agrochemical compositions according to the invention. Genetically modified plants are plants into in which genome a heterologous gene encoding a protein of interest has been stably integrated. The expression "heterologous gene encoding a protein of interest" essentially means  
20 genes which give the transformed plant new agronomic properties, or genes for improving the agronomic quality of the transformed plant.

          The dose of active material usually applied in the treatment according to the present invention is generally and advantageously between 10 and 2000 g/ha,  
25 preferably between 20 and 1500 g/ha for applications in foliar treatment. The dose of active substance applied is generally and advantageously between 1 and 200 g per 100 kg of seed, preferably between 2 and 150 g per 100 kg of seed in the case of seed treatment. It is clearly understood that the doses indicated above are given as illustrative examples of the invention. A person skilled in the art will know how to  
30 adapt the application doses according to the nature of the crop to be treated.

          The compositions according to the present invention may also be used fore the preparation of composition useful to curatively or preventively treat human and animal fungal diseases such as, for example, mycoses, dermatoses, trichophyton  
35 diseases and candidiases or diseases caused by *Aspergillus spp.* or *Candida spp.*, for example *Aspergillus fumigatus* or *Candida albicans* respectively.



The present invention will now be illustrated with the following examples :

**Example 1 : Efficacy against *Pyrenophora teres* of a composition containing N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (Compound 1) and trifloxystrobin.**

The active ingredients tested are prepared by potter homogenisation in a mixture of acetone/tween/water . This suspension is then diluted with water to obtain the desired active material concentration.

Barley plants (Express variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 12°C, are treated at the 1-leaf stage (10 cm tall) by spraying with the aqueous suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material.

After 24 hours, the plants are contaminated by spraying them with an aqueous suspension of *Pyrenophora teres* spores (12,000 spores per ml). The spores are collected from a 12-day-old culture .The contaminated barley plants are incubated for 24 hours at about 20°C and at 100% relative humidity, and then for 12 days at 80% relative humidity.

Grading (% of efficacy) is carried out 12 days after the contamination, in comparison with the control plants.

The following table summarises the results obtained when tested compound 1 and trifloxystrobin alone and in a 1/1 weight ratio mixture.

	Dose (g/ha)	% Efficacy	Synergism (Colby)
Compound 1	4	20	-
	8	50	-
Trifloxystrobin	4	50	-
	8	70	-
Compound 1 + Trifloxystrobin (Ratio 1/1)	4+4	85	+25
	8+8	90	+5

According to the Colby method, a synergistic effect of the mixtures tested has been observed.

**Example 2 : Efficacy against *Alternaria brassicae* (in vitro) of a composition containing N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide (Compound 1) and trifloxystrobin.**

5 The growth of *Alternaria brassicae* was followed on PDA medium containing the compounds alone or in mixtures.

The PDA medium is prepared by mixing 39 grams of PDA (Merck) in 1 liter of demineralized water. The medium is sterilized by autoclave 15 minutes at 121°C. The compounds are dissolved in acetone and added to the medium before pouring it  
10 in Petri dishes. After medium drying, the fungus is inoculated on the medium and allowed to growth at 19°C.

The percentage of efficacy is noted after 14 days of growth (measurement of the radial growth) of the fungus in comparison with a control.

15 The following table summarises the results obtained when tested compound 1 and trifloxystrobin alone and in different weight ratio mixtures.

	Dose (ppm)	% Efficacy	Synergism (Colby)
Compound 1	0.025	34	-
	0.05	45	-
	0.1	53	-
	0.25	76	-
	0.5	90	-
Trifloxystrobin	0.025	45	-
	0.05	44	-
	0.1	45	-
	0.25	53	-
	0.5	60	-
	1	65	-
	2.5	71	-
Compound 1 + Trifloxystrobin (Ratio 1/1)	0.05+0.05	79	+10
	0.1+0.1	93	+19
	0.5+0.5	100	+4
Compound 1 + Trifloxystrobin (Ratio 1/2)	0.05+0.1	85	+15
	0.25+0.5	100	+10
Compound 1 + Trifloxystrobin (Ratio 2/1)	0.1+0.05	97	+23
	0.5+0.25	100	+5
Compound 1 + Trifloxystrobin (Ratio 1/4)	0.025+0.1	75	+11

Compound (Ia) + Trifloxystrobin (Ratio 4/1)	0.1+0.025	100	+26
Compound (Ia) + Trifloxystrobin (Ratio 1/10)	0.05+0.5	96	+18
	0.25+2.5	100	+7

According to the Colby method, a synergistic effect of the mixtures tested has been observed.

5 **Example 3 : Efficacy against *Alternaria brassicae* (in vitro) of a composition containing N-(2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl)-2-trifluoromethylbenzamide (Compound 1) and azoxystrobin.**

10 The growth of *Alternaria brassicae* was followed on PDA medium containing the compounds alone or in mixtures.

The PDA medium is prepared by mixing 39 grams of PDA (Merck) in 1 liter of demineralized water. The medium is sterilized by autoclave 15 minutes at 121°C. The compounds are dissolved in acetone and added to the medium before pouring it in Petri dishes. After medium drying, the fungus is inoculated on the medium and  
15 allowed to growth at 19°C. The percentage of efficacy is noted after 14 days of growth (measurement of the radial growth) of the fungus in comparison with a control.

The following table summarises the results obtained when tested compound 1 and azoxystrobin alone and in different weight ratio mixtures.

20

	Dose (ppm)	% Efficacy	Synergism (Colby)
Compound 1	0.025	34	-
	0.05	45	-
	0.1	53	-
	0.25	76	-
	0.5	90	-
Azoxystrobin	0.025	38	-
	0.05	41	-
	0.1	47	-
	0.25	56	-
	0.5	47	-
	1	48	-
	2.5	49	-

Compound 1 + Azoxystrobin (Ratio 1/1)	0.05+0.05	70	+2
	0.1+0.1	88	+13
	0.5+0.5	100	+5
Compound 1 + Azoxystrobin (Ratio 1/2)	0.05+0.1	76	+5
	0.25+0.5	95	+8
Compound 1 + Azoxystrobin (Ratio 2/1)	0.1+0.05	81	+9
	0.5+0.25	98	+2
Compound (Ia) + Azoxystrobin (Ratio 4/1)	0.1+0.025	100	+29
Compound (Ia) + Azoxystrobin (Ratio 1/10)	0.05+0.5	86	+15
	0.25+2.5	100	+12

According to the Colby method, a synergistic effect of the mixtures tested has been observed.

5 **Example 4 : Efficacy against *Sphaerotheca fuliginea* of a composition containing N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethyl-benzamide (Compound 1) and boscalid.**

10 The active ingredients tested are prepared by potter homogenisation in a mixture of acetone/tween/water . This suspension is then diluted with water to obtain the desired active material concentration.

15 Gherkin plants (Vert petit de Paris variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 20°C/23°C, are treated at the 2-leaves stage by spraying with the aqueous suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material.

After 24 hours, the plants are contaminated by spraying them with an aqueous suspension of *Sphaerotheca fuliginea* spores (100 000 spores per ml). The spores are collected from a contaminated plants .The contaminated gerkin plants are incubated at about 20°C/25°C and at 60/70% relative humidity.

20 Grading (% of efficacy) is carried out 21 days after the contamination, in comparison with the control plants.

The following table summarises the results obtained when tested compound 1 and boscalid alone and in a 1/1 weight ratio mixture.

	Dose (ppm)	% Efficacy	Synergism (Colby)
Compound 1	4	10	-
	8	15	-
	15	25	-
Boscalid	4	0	-
	8	10	-
	15	45	-
Compound 1 + Boscalid (Ratio 1/1)	4+4	40	+30
	8+8	70	+47
	15+15	95	+36

According to the Colby method, a synergistic effect of the mixtures tested has been observed.

**Example 5 : Efficacy against *Sphaerotheca fuliginea* of a composition containing N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethyl-benzamide (Compound 1) and N-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoro-methyl)-1-methyl-1H-pyrazole-4-carboxamide (Compound A).**

The active ingredients tested are prepared by potter homogenisation in a mixture of acetone/tween/water . This suspension is then diluted with water to obtain the desired active material concentration.

Gherkin plants (Vert petit de Paris variety) in starter cups, sown on a 50/50 peat soil-pozzolana substrate and grown at 20°C/23°C, are treated at the 2-leaves stage by spraying with the aqueous suspension described above. Plants, used as controls, are treated with an aqueous solution not containing the active material.

After 24 hours, the plants are contaminated by spraying them with an aqueous suspension of *Sphaerotheca fuliginea* spores (100 000 spores per ml). The spores are collected from a contaminated plants .The contaminated gherkin plants are incubated at about 20°C/25°C and at 60/70% relative humidity.

Grading (% of efficacy) is carried out 21 days after the contamination, in comparison with the control plants.

The following table summarises the results obtained when tested compound 1 and boscalid alone and in a 1/1 weight ratio mixture.


	Dose (ppm)	% Efficacy	Synergism (Colby)
Compound 1	8	75	-
Compound A	8	60	-
Compound 1 + Compound A (Ratio 1/1)	8+8	95	+5

According to the Colby method, a synergistic effect of the mixture tested has been observed.

CLAIMS

1. A composition comprising :

5 a) a pyridylethylbenzamide derivative of general formula (I)



in which :

- p is an integer equal to 1, 2, 3 or 4;
- q is an integer equal to 1, 2, 3, 4 or 5;
- each substituent X is chosen, independently of the others, as being halogen, alkyl or

10 haloalkyl;

- each substituent Y is chosen, independently of the others, as being halogen, alkyl, alkenyl, alkynyl, haloalkyl, alkoxy, amino, phenoxy, alkylthio, dialkylamino, acyl, cyano, ester, hydroxy, aminoalkyl, benzyl, haloalkoxy, halosulphonyl, halothioalkyl, alkoxyalkenyl, alkylsulphonamide, nitro, alkylsulphonyl, phenylsulphonyl or

15 benzylsulphonyl;

as to the N-oxides of 2-pyridine thereof;

and

b) a compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms;

20 in a (a) / (b) weight ratio of from 0.01 to 20.

2. A composition according to claim 1, characterised in that p is 2.

3. A composition according to claim 1 or 2, characterised in that q is or 2.

25

4. A composition according to any of the claims 1 to 3, characterised in that X is chosen, independently of the others, as being halogen or haloalkyl.

5. A composition according to any of the claims 1 to 4, characterised in that X is

30 chosen independently of the others, as being a chlorine atom or a trifluoromethyl group.

6. A composition according to any of the claims 1 to 5, characterised in that Y is chosen, independently of the others, as being halogen or haloalkyl.
- 5 7. A composition according to any of the claims 1 to 6, characterised in that Y is chosen, independently of the others, as being a chlorine atom or a trifluoromethyl group.
8. A composition according to any of the claims 1 to 7, characterised in that the  
10 compound of general formula (I) is :  
- N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide;  
- N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-iodobenzamide; or  
- N-{2-[3,5-dichloro-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide .
- 15 9. A composition according to claim 8, characterised in that the compound of general formula (I) is N-{2-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]ethyl}-2-trifluoromethylbenzamide.
- 10 10. A composition according to any of the claims 1 to 9, characterised in that the compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms is a compound capable of inhibiting reduced nicotinamide-adenine dinucleotide dehydrogenase in phytopathogenic fungal organisms.
- 25 11. A composition according to claim 10, characterised in that the compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms is diflumetorin.
- 30 12. A composition according to any of the claims 1 to 9, characterised in that the compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms is a compound capable of inhibiting succinate dehydrogenase in phytopathogenic fungal organisms.
- 35 13. A composition according to claim 12 characterised in that the compound capable of inhibiting the transport of electrons of the respiratory chain of succinate dehydrogenase in phytopathogenic fungal organisms is N-[2-(1,3-dimethyl-butyl)-



phenyl]-5-fluoro-1,3-dimethyl-1*H*-pyrazole-4-carboxamide, *N*-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoro-methyl)-1-methyl-1*H*-pyrazole-4-carboxamide, *N*-[2-(1,3-dimethylbutyl)-thiophen-3-yl] 1-methyl-3-(trifluoromethyl)-1*H*-pyrazole-4-carboxamide, benodanil, carboxin, fenfuram, flutolanil, furametpyr, mepronil,  
 5 boscalid, oxycarboxin or thifluzamide.

14. A composition according to any of the claims 1 to 9, characterised in that the compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms is a compound capable of inhibiting  
 10 mitochondrial ubiquinol:ferricytochrome-c oxidoreductase in phytopathogenic fungal organisms.

15. A composition according to claim 14, characterised in that the compound capable of inhibiting the transport of electrons of the respiratory chain of  
 15 mitochondrial ubiquinol:ferricytochrome-c oxidoreductase in phytopathogenic fungal organisms is a strobilurin derivative, cyazofamid, fenamidone or famoxadone.

16. A composition according to claim 15, characterised in that the strobilurin derivative is azoxystrobin, dimoxystrobin, fluoxastrobin, kresoxim-methyl,  
 20 metominostrobin, trifloxystrobin, pyraclostrobin, picoxystrobin or 2-{2-[6-(3-chloro-2-methylphenoxy)-5-fluoro-pyrimidin-4-yloxy]-phenyl}2-methoxyimino-*N*-methylacetamide.

17. A composition according to any one of the claims 1 to 16 further comprising  
 25 a fungicidal compound (c).

18. A composition according to claim 17, characterised in that the fungicidal compound (c) is selected from captane, folpet, dodine, propineb, mancozeb, thiram, tolylfluanid, iminoctadine, dithianon, copper hydroxide, copper octanoate, copper  
 30 oxychloride, copper sulfate, fosetyl-Al, phosphorous acid, cymoxanil, iprovalicarb, benthiavalicarb, chlorotalonil, propamocarb, tebuconazole and spiroxamine.

19. A composition according to any one of the claims 1 to 18, characterised in that it further comprises an agriculturally acceptable support, carrier, filler and/or  
 35 surfactant.

20. A method for preventively or curatively controlling phytopathogenic fungi of crops, characterised in that an effective and non-phytotoxic amount of a composition according to any one of the claims 1 to 19 is applied to the seed, the plant and/or to the fruit of the plant or to the soil in which the plant is growing or in which it is desired to  
5 grow.

### ABSTRACT

5 A composition comprising at least a pyridylethylbenzamide derivative of general formula (I) (a) and a compound capable of inhibiting the transport of electrons of the respiratory chain in phytopathogenic fungal organisms (b) in a (a) / (b) weight ratio of from 0.01 to 20.

A composition further comprising an additional fungicidal compound.

10

A method for preventively or curatively combating the phytopathogenic fungi of crops by using this composition.